

Level and Distribution of Heavy Metals in Miri River, Miri City, Malaysia: A Case Study

Amit Kumar Maharjan^{1*}, Dick Rong En Wong^{2*}, Rubiyatno Rubiyatno³

¹Interdisciplinary Centre for River Basin Environment, University of Yamanashi, Japan

²Faculty of Engineering and Science, Curtin University, CDT250 Miri 98009, Malaysia

³Integrated Graduate School of Medicine, Engineering, and Agricultural Sciences, University of Yamanashi, Japan

*Correspondence: amit_kmr@hotmail.com; 700014455@student.curtin.edu.my

SUBMITTED: date; REVISED: 27 August 2021; ACCEPTED: 2 September 2021

ABSTRACT: The heavy metals pollution in the water resource has become a serious and hazardous environmental problem all over the world because of non-biodegradability, emanating from multiple sources, easy accumulation, and biological toxicity. This research was carried out to study the level and distribution of heavy metals at different sampling locations (upstream, midstream, and downstream), at different depths (0.5 m and 1.5 m from surface water level), and during low tide and high tide conditions in the Miri River of Miri City in Malaysia. The river water samples were collected and analyzed for Ca, Mg, Cu, Fe, Mn, Ni, Pb, and Zn by flame atomic absorption spectrophotometer. Concentration of Ca was found to be the highest in the Miri River, followed by Mg and Fe, and with traces of Cu, Mn, Ni, Pb, and Zn. Increment in the concentration of heavy metals, such as Cu, Mg, and Ni, was observed while flowing from upstream to downstream of the Miri River. Concentration of heavy metals, such as Ca, Mg, Cu, and Zn, were clearly lower at 1.5 m depth than at 0.5 m depth. High tides in the river decreased the concentration of heavy metals, such as Ca, Cu, Mn, and Ni, than during low tides. From this research, it gets clear that using the Miri River water for domestic and recreational purposes, washing, and fishing is detrimental to human health and the environment.

KEYWORDS: heavy metals; Miri River; river water quality; depth; tides

1. Introduction

Water resources are important natural resource for life on earth. Water is irrefutable resource for domestic purposes, agriculture (e.g. irrigation, animal husbandry, etc.), various industries, energy productions, and provide many ecosystem services (e.g. tourism, recreational activities, etc.). However, various natural and anthropogenic activities degrade the quality of water resources and impair their use [1,2]. Water pollution has gained a serious concern, especially in developing countries due to direct discharge of wastewater into rivers and lakes, without treatment [3–5]. At present, water quality is one of the key problems in the aquatic environment. Several point and non-point sources are contributing the river water pollutions [6]. The direct discharge of wastes and industrial by-products associated with toxic compounds, without proper treatment, into water resources represents an ongoing environmental problem

due to their possible impact on aquatic environment or river ecosystem and a potential effect on public health [7]. The pollution of water resources directly affects the quality and quantity of water for irrigation and drinking water and increases the financial burden to treat or make it useable [6].

The trace metal pollution in water resource has become a serious and hazardous environmental problem all over the world because of non-biodegradability, emanating from multiple sources, easy accumulation, and biological toxicity [8]. Trace metals are the dense metallic component, naturally occurring on the earth's crust, which are toxic to both aquatic and human life, due to their persistent and non-biodegradable characteristics with the ability to cause deleterious effects on living organisms [9]. Various solid and liquid wastes are generated as the by-products of manufacturing processes, which contains toxic chemicals such as chromium salts, sulfides and other substances including heavy toxic trace metals [10]. Heavy metal pollution can occur through natural processes such as geological weathering, direct atmospheric deposition, etc., and anthropogenic activities such as agricultural, municipal, residential or industrial waste products [11]. A major source of energy for industries is coal combustion, which is one of the most important anthropogenic emission sources of trace elements and an important source of a number of metals [12].

Continuous accumulation of metals in water bodies can amplify the adverse effects on aquatic life and water ecosystem. Human body requires some metals such as zinc, copper, iron, manganese, and cobalt for their development in very low concentrations, however, they may be toxic if ingested at higher concentrations [13]. Mostly, heavy metals enter human body through ingestion (the consumption of contaminated drinking water or food) and may result in cardiovascular disorders, neuronal damage, renal injuries, and risk of cancer and diabetes [14,15]. Trace metals can cause deoxyribonucleic acid (DNA) damage and lesions through the formation of reactive oxygen species in tissues [16]. Moreover, the irrigation with the heavy metals contaminated water degrades the soil quality and affects overall agricultural system including growth and yield of plants, consequently the animals consuming the products [17–19].

Around 98% of the total water used in Malaysia is from rivers [20]. According to a report published by Department of Energy, Malaysia, in 2017, out of 477 rivers being monitored, about 11% of the rivers were found as polluted and about 43% as slightly polluted. Miri River is the major surface water source, flowing through the Miri city. Miri River starts from inland, flows through the Miri city, curving at Lutong town, and ends out to South China Sea. Miri River contributes 97% of the total water use, mainly irrigation and domestic needs of the Miri city, all around a year [21]. Hence, it is important to study the level and distribution of heavy metals in the water of Miri River. The objective of this research was to study the level of heavy metals in the Miri River of Miri City, Sarawak, Malaysia and investigate the distribution of heavy metals at different sampling points in the river, depth of water in the river, and effects of tide on the concentration of the heavy metals.

2. Materials and Methods

2.1. Description of the Study Area

Miri is a coastal city in northeastern Sarawak, Malaysia, located near the border of Brunei, on the island of Borneo. The city is the capital of Miri District of the Miri Division and the second

largest city in Sarawak. Population of the city is increasing every year, which emerged the city as an industrial district and many factories are built in the city, along the river. The establishment of these factories often entailed significant environmental footprints, due to the direct disposal of biological and chemical wastes of the factories in the Miri River. One of the major pollutant in the river is heavy metals. The heavy metals that pervaded into the river would pollute the river which resulted in a poor water quality condition.

2.2. Water Sampling

Water sampling locations were determined at the upstream, midstream (about 7.5 km river length downstream from the upstream locations), and downstream (about 7.4 km river length downstream from the midstream locations) of the Miri River and samples were collected from 2 points in each locations by grab method. Exact location and description of the sampling points are presented in Table 1. Water samples were collected from 0.5 m and 1.5 m depth from the surface water level of each sampling points, to check the variation in concentrations at different depths. Also, the water samples were collected during low tide and high tide period, to understand the effect of tides in the variation of concentrations.

Table 1. Location (coordinates) and description of the sampling points.

	Coordinate of point 1	Coordinate of point 2	Description
Upstream	4.412236,114.019651	4.411019,114.020787	Heavy industrial and residential area
Midstream	4.468002,114.009296	4.467889,114.012893	Commercial area
Downstream	4.400356,113.986662	4.399246,113.986053	Fishery and industrial area

2.3. Preparation and Analysis of Samples

The collected water samples were carried to laboratory, acidified with 20% nitric acid, digested, filtered, and analyzed for eight different heavy metals, namely lead (Pb), iron (Fe), manganese (Mn), copper (Cu), nickel (Ni), magnesium (Mg), calcium (Ca), and zinc (Zn). The samples were analyzed by using Flame Atomic Absorption Spectrophotometer (AA400, Perkin Elmer), following Standard Methods for the Examination of Water and Wastewater [22]. Various concentrations of standard solution of same heavy metal was prepared by dilution, and was used to calibrate and analyze the river water samples. Different wavelength of source lamp and different type of lamp was used for the analysis of each heavy metal.

3. Results and Discussion

The results of the study are presented in this section. Result or concentration of each heavy metal at different sampling points (upstream [U/S], midstream [M/S], and downstream [D/S]) of the Miri river, at different depth (0.5 m and 1.5 m depth), and the effect of tides (low tide [LT] and high tide [HT]) on the variation of concentration are presented and discussed individually for each metal.

3.1. Calcium (Ca)

Concentration of Ca at upstream, midstream, and downstream of the Miri River, at 0.5 m depth and 1.5 m depth, and variation in concentration at low tide and high tide conditions are presented in Figure 1. The Ca concentration in the Miri River was found to be widely varied in the range of 113 to 1,545 mg/L. The Ca concentration was found to be fluctuating in the

upstream, midstream, and downstream sampling locations and different sampling depths. In general, the concentration of Ca in rivers are low, but concentration can be high due to flow through the lime areas or dumping of the wastes containing Ca. Human activities (domestic and industrial) in the vicinity of the Miri River might have contributed to the higher concentration of Ca. Sources of Ca can be some industrial waste waters from processing, in which the acids are neutralized by lime or limestone [23]. Though, there are no health based value for the concentration of Ca in WHO guidelines for drinking water, the taste threshold for the Ca ion is in the range of 100-300 mg/L [13]. Also, there remains the possibility of scale deposition in the treatment works, distribution system and pipeworks of the water containing high Ca concentrations. There is no limit provided in the guideline for Ca concentration in the National Water Quality Standards (NWQS) for Malaysia [24], but the water hardness should be paid attention, before using the river water for any type of use.




Figure 1. Calcium (Ca) concentration at different location of the Miri River, at different depth, and variation on concentration due to tidal effect. The plots present mean values of 2 sampling points at each location and the error bars correspond to S.D. ($n = 2$).

3.2. Magnesium (Mg)




Figure 2. Magnesium (Mg) concentration at different location of the Miri River, at different depth, and variation on concentration due to tidal effect. The plots present mean values of 2 sampling points at each location and the error bars correspond to S.D. ($n = 2$).

Concentration of Mg at upstream, midstream, and downstream of the Miri River, at 0.5 m depth and 1.5 m depth, and variation in concentration at low tide and high tide conditions are presented in Figure 2. The Mg concentration in the Miri River ranged from 29 to 217 mg/L. There are no health based value for Mg concentration in WHO guidelines for drinking water,

but it can increase the hardness of water, limiting the usability of the water. The Mg concentration was found to be fluctuating in the upstream, midstream, and downstream sampling locations and different sampling depths. However, the Mg concentration was found to be increasing from upstream to downstream, which might be due to direct discharge of wastes to the river at the downstream. The Mg concentration was lower during high tide than during low tide. It might be due to the dilution of Mg in higher volume of water, during high tide condition.

3.3. Copper (Cu)

Cu is an important trace element (recommended daily allowance of 2.0-3.0 mg/day for human) to several enzymes involved in antioxidant response, membrane and DNA integrity, and adenosine triphosphate production [25,26]. Accumulation of Cu may cause protein damage and cell injury in bacteria [26,27] and high levels of Cu exposure may induce toxicity and the development of copper resistance in various pathogenic bacteria [28,29]. Concentration of Cu at upstream, midstream, and downstream of the Miri River, at 0.5 m depth and 1.5 m depth, and variation in concentration at low tide and high tide conditions are presented in Figure 3. The Cu concentration in the Miri River ranged from 0.03 to 0.13 mg/L.

Naturally occurring concentration of Cu in the freshwater systems had been reported to be from 0.20 to 30 $\mu\text{g/L}$ [30]. It was observed that the Cu concentrations increased from upstream to downstream and the concentration was lower during high tide than that during low tide. Human activities and the establishment of industries near the Miri River might have contributed as the anthropogenic sources for increase in concentration of Cu. According to the NWQS for Malaysia [24], the Miri River water generally did not conform to the established standard for drinking water and recreations involving body contact. The water might also impact fishery as the water was not suitable for sensitive and very sensitive aquatic species. The water could be used for irrigation as well as deemed acceptable for common aquatic species.




Figure 3. Copper (Cu) concentration at different location of the Miri River, at different depth, and variation on concentration due to tidal effect. The plots present mean values of 2 sampling points at each location and the error bars correspond to S.D. ($n = 2$).

3.4. Iron (Fe)

Fe is one of the most abundant metals in Earth's crust and essential element in human nutrition. No health based value has been provided for concentration of Fe in WHO guidelines for drinking water [13], but preferred below 0.3 mg/L, as the taste of Fe is unnoticeable till that

value. Higher concentrations of Fe affects the taste of drinking water and stains the clothes during laundry. Concentration of Fe at upstream, midstream, and downstream of the Miri River, at 0.5 m depth and 1.5 m depth, and variation in concentration at low tide and high tide conditions are presented in Figure 4. The Fe concentration in the Miri River ranged from 1.45 to 7.63 mg/L. Fe concentration in the upstream (> 5 mg/L) was significantly higher than that in the midstream and downstream. According to the NWQS for Malaysia [24], the Miri River water at the upstream was not suitable even for the irrigation purpose. Higher concentration of Fe at the upstream and lower concentration of Fe at the midstream and downstream might be due to settling of the Fe-sediments in the river-bed during flow or the discharge of Fe wastes nearby the upstream sampling points.




Figure 4. Iron (Fe) concentration at different location of the Miri River, at different depth, and variation on concentration due to tidal effect. The plots present mean values of 2 sampling points at each location and the error bars correspond to S.D. ($n = 2$).

3.5. Manganese (Mn)

Mn is also an abundant metals in Earth's crust and commonly occurring with Fe. Concentration of Mn in a naturally occurring surface waters ranged from 1 to 200 $\mu\text{g/L}$. A health based value of 0.4 mg/L of Mn concentration has been considered in the WHO guidelines for drinking water [13]. If the concentration of Mn is below 0.1 mg/L, then both guidelines for organoleptic properties and health guidelines can be met [31]. Concentration of Mn at upstream, midstream, and downstream of the Miri River, at 0.5 m depth and 1.5 m depth, and variation in concentration at low tide and high tide conditions are presented in Figure 5. The Mn concentration in the Miri River ranged from 0.03 to 0.48 mg/L. Concentration of Mn at the upstream and midstream were less than 0.11 mg/L (at both sampling depths and tidal conditions), whereas the concentration was higher in downstream. It might be due to the disposal of wastes containing Mn or Mn compounds in the downstream. According to NWQS for Malaysia [24], the Miri River water at upstream and downstream can be used for drinking and recreational purposes, after proper treatment, from the view point of concentration of Mn, however, the Miri River water at downstream was not fit for drinking and recreational purposes as well as could cause adverse impacts to fishery industries, but can be used for irrigation. The Miri River water at upstream and midstream can also be used for fisheries of sensitive and common aquatic species. This might be the reason for establishment of many fisheries at the downstream of Miri River.




Figure 5. Manganese (Mn) concentration at different location of the Miri River, at different depth, and variation on concentration due to tidal effect. The plots present mean values of 2 sampling points at each location and the error bars correspond to S.D. ($n = 2$).

3.6. Nickel (Ni)

The maximum recommended value of Ni based on human health criteria for drinking water is 0.07 mg/L. Higher concentrations of Ni in water causes several health risks to plants and animals [13]. Concentration of Ni at upstream, midstream, and downstream of the Miri River, at 0.5 m depth and 1.5 m depth, and variation in concentration at low tide and high tide conditions are presented in Figure 6. The Ni concentration in the Miri River ranged from 0.12 to 0.26 mg/L. The concentration of Ni in the Miri River water was higher than the guidelines value. It might be due to the release of Ni and Ni compounds from natural or industrial Ni wastes in the river bank. Concentration of Ni increased in the downstream than in the upstream. It might be due to the disposal of industrial wastes containing Ni in the downstream, without proper treatment. In general, the concentration of Ni was lower in the higher depth (i.e., 1.5 m) than that in the lower depth (i.e., 0.5 m), and also in high tide condition than that in low tide condition. It might be due to the dilution of Ni in the higher volume of river water during high tide conditions. According to NWQS for Malaysia [24], the Miri River water was found to be unfit for drinking and recreational activities involving body contact, from the aspect of Ni contamination. However, the concentration of Ni in water at the upstream and midstream of Miri River were less than 0.2 mg/L, which indicates that the river water at the upstream and downstream can be used for irrigation purpose.




Figure 6. Nickel (Ni) concentration at different location of the Miri River, at different depth, and variation on concentration due to tidal effect. The plots present mean values of 2 sampling points at each location and the error bars correspond to S.D. ($n = 2$).

3.7. Lead (Pb)

Pb is one of the commonest heavy metal. Pb has been recognized and highlighted as one of the most dangerous environmental poisons [32]. The most recent guideline value for Pb in drinking water from the WHO is 10 µg/L but is no longer a health based value and has been designated provisional [13], due to understanding of harmful impacts of Pb on health. Concentration of Pb at upstream, midstream, and downstream of the Miri River, at 0.5 m depth and 1.5 m depth, and variation in concentration at low tide and high tide conditions are presented in Figure 7. The Pb concentration in the Miri River fluctuated from not detected to 0.35 mg/L. The Pb concentration in the normal rivers range from 0.003 to 0.03 mg/L. This indicates that the Miri River is polluted in the aspect of Pb concentration too. Though any particular trend of variation in Pb concentration at the upstream, midstream, and downstream sampling locations, depths, and tidal conditions was not obtained, it can be observed that the Miri River water was not suitable for drinking, and recreational activities involving body contact. According to NWQS for Malaysia [24], the Miri River water could also impact the fisheries or aquatic species of sensitive nature, and recommended for irrigation purpose only. Pb usually gets into water bodies due to leaching from lead-containing service lines, lead solder, and brass fittings, particularly in corrosive waters. There might be similar industries along the Miri River and discharging their waste without proper treatment.




Figure 7. Lead (Pb) concentration at different location of the Miri River, at different depth, and variation on concentration due to tidal effect. The plots present mean values of 2 sampling points at each location and the error bars correspond to S.D. ($n = 2$).

3.8. Zinc (Zn)

Zn is an essential element required by human for optimum growth and development, but the presence of high amount of Zn in water may cause a bitter taste and opalescence in water. Though there is no health based value for Zn concentration in WHO guidelines for drinking water, but the preferred limit is 3 mg/L [13]. Concentration of Zn at the upstream, midstream, and downstream of the Miri River, at 0.5 m depth and 1.5 m depth, and variation in concentration at low tide and high tide conditions are presented in Figure 8. The Zn concentration in the Miri River ranged from 0.04 to 0.18 mg/L. This indicates that the Miri River water is fairly acceptable for every uses, in the aspect of Zn concentration [24]. However, no any particular trend of variation in Zn concentration at the upstream, midstream, and downstream sampling locations, depths, and tidal conditions was obtained.




Figure 8. Zinc (Zn) concentration at different location of the Miri River, at different depth, and variation on concentration due to tidal effect. The plots present mean values of 2 sampling points at each location and the error bars correspond to S.D. ($n = 2$).

3.9. Heavy Metals in the Miri River

Levels and distributions in concentration of eight heavy metals, namely Ca, Mg, Cu, Fe, Mn, Ni, Pb, and Zn, were studied at the upstream, midstream, and downstream sampling locations, at 0.5 m depth and 1.5 m depth, and at high tide and low tide conditions in the Miri River. Concentrations of Ca was found to be the highest in the Miri River, followed by Mg and Fe. Concentration of other metals (Cu, Mn, Ni, Pb, and Zn) were in traces but few of them were higher than the permissible limits of drinking water and some even can not be used for recreational as well as fisheries (discussed individually in previous section). Previous study by Wogu and Okaka in the Warri River in Nigeria [33], which receives the industrial, agricultural, and urban sewage, showed that the concentration of Cd, Cr, Mn, and Ni were higher than the standard level, and the water was harmful to the public health and hygiene.

The results showed the increase in concentration of heavy metals from upstream to downstream, especially, for the Cu, Mg, and Ni. The results of the analysis of water and sediment samples in upstream and downstream of the Tembi River in Iran [34] also revealed that the mean concentration of several heavy metals in the downstream river water was significantly higher than that in the upstream river water. As the water flows downstream of the river, the dissolved heavy metals and sediments in upstream are also transported to downstream. Physicochemical characteristics of water might also affect the precipitation of the heavy metals in the sediments [35] and contribute to the increase in concentration of heavy metals in the downstream. In addition to this, the human activities and direct disposal of industrial wastes in the vicinity of the Miri River in the midstream and downstream might have increased the concentrations of heavy metals in downstream of the Miri River.

Anomalies in the variation of concentration of some heavy metals in upstream and downstream existed. For example, the concentration of Fe was higher in the upstream than in the downstream. It could be due to the sampling locations in the upstream, being near to the discharge points of domestic and industrial wastes. Similar phenomenon was observed in a study at Yamuna River in India [36]. The Fe concentration increased many folds in the downstream, because of several iron works or related industries in the downstream area. Concentration of heavy metals, such as Ca, Mg, Cu, and Zn, were clearly lower at 1.5 m depth than at 0.5 m depth, while fluctuating results for other metals. In general, concentration of the heavy metals increased with depth of sampling in a study of depth variability of heavy metal

concentrations in the water of Firiza-Strimtori Lake, NW of Romania [37]. It might be due to the emanation of the metal ions into the water as an endogenic source of contamination based on the alteration in the re-dox potential of sediments. Anomalies in the result of other metals, might be due to the variation in pH, re-dox potential, and river flow characteristics (e.g., flow rate) [38].

Concentration of heavy metals, such as Ca, Cu, Mn, and Ni, were observed to be lower in high tide conditions than that in low tide condition, whereas no clear trend was obtained in remaining metals. The decrease in concentration at high tide condition can be directly linked with the dilution of the metals in huge volume of water during high tide, in comparison to water volume during low tide. Tidal currents can affect the shifting of pollutants along the estuary, moving upstream during high tide and downstream during low tide [38], which could be one reason of anomalies in the variation of concentration of heavy metals in upstream and downstream.

A part from these parameters, the level and distribution of heavy metals depend on the surrounding environmental factors and several physico-chemical characteristics [39,40]. Fine-grained sediments has higher surface area for adsorption and ionic attraction, so, the concentration of heavy metals, generally, increased with smaller particle size of sediments [41,42]. Moreover, the transportation of sediments, along with the heavy metals, to the downstream of the river, were also higher with the fine-grained sediments [42]. The pH, dissolved oxygen, temperature, and flow rate of the river water also affects the release of heavy metals from sediments [43]. Higher hydrogen ion concentration at lower pH, tends to occupy more adsorption sites in sediments, resulting in easier precipitation of soluble and carbonate-bound heavy metals. Heavy metals can exist in the combined form with organic matters, which will oxidize with higher dissolved oxygen in the water, and release heavy metals from organic matters in water. Higher temperature can accelerate the release of the heavy metals in water from water-soluble fractions, carbonate fractions, and exchangeable fraction from the sediments. With the higher flow rate, there remains the possibility of higher dissolved oxygen and the physical disturbance in the sediments, which can promote the oxidation/reaction to release the heavy metals in water. However, these parameters were not considered in this study.

4. Conclusions

The level of heavy metals at different sampling locations (upstream, midstream, and downstream), at different depths (0.5 m and 1.5 m from surface water level), and at low tide and high tide conditions of the Miri River were studied. Concentration of Ca was found to be the highest in the Miri River, followed by Mg and Fe. Concentration of other metals (Cu, Mn, Ni, Pb, and Zn) were in traces but few of them were higher than the permissible limits of drinking water and some even can not be used for recreational purposes involving body contact as well as fisheries. Increment in the concentration of heavy metals, such as Cu, Mg, and Ni, was observed from upstream to downstream of the Miri River, whereas, some anomalies existed. Concentration of heavy metals, such as Ca, Mg, Cu, and Zn, were clearly lower at 1.5 m depth than at 0.5 m depth, while results fluctuated for other metals. Concentration of heavy metals, such as Ca, Cu, Mn, and Ni, were observed to be lower during high tide condition than that during low tide condition, whereas no clear trend was obtained in remaining metals. In general, the Miri River water was found to be polluted with heavy metals, at different concentration levels. According to the NWQS for Malaysia, Miri River water can be considered

as suitable for the irrigation purpose only. It is recommended for integrated and comprehensive study of the heavy metal concentrations in the Miri River, considering all the affecting parameters, to better understand the distribution of heavy metals and sources of contaminations.

Acknowledgments

The authors thank Curtin University Malaysia for facilitating this study. Collaboration from Yamanashi University is highly appreciated.

Competing Interest

All authors declare no competing interests.

References

- [1] Carpenter S.R.; Caraco N.F.; Correll D.L.; Howarth R.W.; Sharpley A.N.; Smith V.H. (1998). Nonpoint pollution of surface waters with phosphorus and nitrogen. *Ecological Application*, 8, 559–568. [https://doi.org/10.1890/1051-0761\(1998\)008\[0559:NPOSWW\]2.0.CO;2](https://doi.org/10.1890/1051-0761(1998)008[0559:NPOSWW]2.0.CO;2).
- [2] Demirak A.; Yilmaz F.; Levent Tuna A.; Ozdemir N. (2006). Heavy metals in water, sediment and tissues of *Leuciscus cephalus* from a stream in southwestern Turkey. *Chemosphere*, 63, 1451–1458. <https://doi.org/10.1016/j.chemosphere.2005.09.033>.
- [3] Unesco WWAP (2017). Wastewater: The untapped resource, Paris. <http://unesdoc.unesco.org/images/0024/002471/247153e.pdf>.
- [4] Abbaspour S. (2011) Water quality in developing countries, South Asia, South Africa, Water quality management and activities that cause water pollution. In International Conference on Environmental and Agriculture Engineering, IACSIT Press, Singapore; pp. 94–102.
- [5] Afroz R.; Masud M.M.; Akhtar R.; Duasa J.B. (2014). Water pollution: Challenges and future direction for water resource management policies in Malaysia. *Environmental Urban ASIA*, 5, 63–81. <https://doi.org/10.1177/0975425314521544>.
- [6] Arefin M.A.; Mallik A. (2017). Sources and causes of water pollution in Bangladesh: A technical overview. *Bibechana*, 15, 97–112. <https://doi.org/10.3126/bibechana.v15i0.18688>.
- [7] Abbas Alkarkhi F.M.; Ismail N.; Easa A.M. (2008). Assessment of arsenic and heavy metal contents in cockles (*Anadara granosa*) using multivariate statistical techniques. *Journal of Hazardous Materieals*, 150, 783–789. <https://doi.org/10.1016/j.jhazmat.2007.05.035>.
- [8] Fan H.; Chen S.; Li Z.; Liu P.; Xu C.; Yang X. (2020). Assessment of heavy metals in water, sediment and shellfish organisms in typical areas of the Yangtze River Estuary, China. *Marine Pollution Bulletin*, 151,. <https://doi.org/10.1016/j.marpolbul.2019.110864>.
- [9] Strungaru S.A.; Nicoara M.; Teodosiu C.; Baltag E.; Ciobanu C.; Plavan G. (2018). Patterns of toxic metals bioaccumulation in a cross-border freshwater reservoir. *Chemosphere*, 207, 192–202. <https://doi.org/10.1016/j.chemosphere.2018.05.079>.
- [10] Tariq S.R.; Shah M.H.; Shaheen N.; Khaliq A.; Manzoor S.; Jaffar M. (2006). Multivariate analysis of trace metal levels in tannery effluents in relation to soil and water: A case study from Peshawar, Pakistan. *Journal of Environmental Management*, 79, 20–29. <https://doi.org/10.1016/j.jenvman.2005.05.009>.
- [11] Dawson E.J.; Macklin M.G. (1998). Speciation of heavy metals in floodplain and flood sediments: a reconnaissance survey of the Aire Valley, West Yorkshire, Great Britain. *Environmental Geochemistry and Health*, 20, 67–76. <https://doi.org/10.1023/A:1006541724394>.
- [12] Wagner A.; Boman J. (2003). Biomonitoring of trace elements in muscle and liver tissue of freshwater fish. *Spectrochimica Acta - Part B Atomic Spectroscopy*, 58, 2215–2226. <https://doi.org/10.1016/j.sab.2003.05.003>.
- [13] WHO (2011). Guidelines for drinking-water quality, 4th ed., World Health Organization, Geneva, Switzerland.
- [14] Martin S.; Griswold W. (2009). Human health effects of heavy metals. *Environmental Science and Technology Briefs from Citizens*, 15, 1–6.

- [15] Rehman K.; Fatima F.; Waheed I.; Akash M.S.H. (2018). Prevalence of exposure of heavy metals and their impact on health consequences. *Journal of Cellular Biochemistry*, 119, 157–184. <https://doi.org/10.1002/jcb.26234>.
- [16] Kim H.S.; Kim Y.J.; Seo Y.R. (2015) An overview of carcinogenic heavy metal: Molecular toxicity mechanism and prevention. *Journal of Cancer Prevention*, 20, 232–240. <https://doi.org/10.15430/jcp.2015.20.4.232>.
- [17] Tutic A.; Novakovic S.; Lutovac M.; Biocanin R.; Ketin S.; Omerovic N. (2015). The heavy metals in agrosystems and impact on health and quality of life. *Macedonian Journal of Medical Sciences*, 3, 345–355. <https://doi.org/10.3889/oamjms.2015.048>.
- [18] Chaoua S.; Boussaa S.; El Gharmali A.; Boumezzough A. (2019). Impact of irrigation with wastewater on accumulation of heavy metals in soil and crops in the region of Marrakech in Morocco. *Journal of Saudi Society of Agricultural Science*, 18, 429–436. <https://doi.org/10.1016/j.jssas.2018.02.003>.
- [19] Sandeep G.; Vijayalatha K.R.; Anitha T. (2019). Heavy metals and its impact in vegetable crops. *International Journal of Chemical Studies*, 7, 1612–1621.
- [20] Huang Y.F.; Ang S.Y.; Lee K.M.; Lee T.S. (2015). Quality of water resources in Malaysia, In *Research and Practices in Water Quality*, pp. 65–94. <https://doi.org/10.5772/58969>.
- [21] Prasanna M. V.; Praveena S.M.; Chidambaram S.; Nagarajan R.; Elayaraja A. (2012). Evaluation of water quality pollution indices for heavy metal contamination monitoring: A case study from Curtin Lake, Miri City, East Malaysia. *Environ. Earth Sci.*, 67, 1987–2001. <https://doi.org/10.1007/s12665-012-1639-6>.
- [22] Rice, E.W.; Baird, R.B.; Eaton, A.D. (1999). *Standard methods for the examination of water and wastewater*, 20th ed.; American Public Health Association, American Water Works Association, Water Environment Federation, Washington DC, USA.
- [23] Awa, S.H.; Hadibarata, T. (2020). Removal of Heavy Metals in Contaminated Soil by Phytoremediation Mechanism: a Review. *Water, Air, & Soil Pollution*, 231, 46. <https://doi.org/10.1007/s11270-020-4426-0>.
- [24] Ministry of Natural Resources and Environment Malaysia (2014). National water quality standards for Malaysia. http://www.wepa-db.net/policies/law/malaysia/eq_surface.htm.
- [25] Nargund S.; Qiu J.; Goudar C.T. (2015). Elucidating the role of copper in CHO cell energy metabolism using 13C metabolic flux analysis. *Biotechnology Progress*, 31, 1179–1186. <https://doi.org/10.1002/btpr.2131>.
- [26] Dalecki A.G.; Crawford C.L.; Wolschendorf F. (2017). *Copper and antibiotics: Discovery, modes of action, and opportunities for medicinal applications*, 1st ed., Elsevier Ltd. <https://doi.org/10.1016/bs.ampbs.2017.01.007>.
- [27] Pontin K.P.; Borges K.A.; Furian T.Q.; Carvalho D.; Wilsmann D.E.; Cardoso H.R.P.; Alves A.K.; Chitolina G.Z.; Salle C.T.P.; Moraes H.L. de S.; do Nascimento V.P. (2021). Antimicrobial activity of copper surfaces against biofilm formation by *Salmonella Enteritidis* and its potential application in the poultry industry. *Food Microbiology*, 94, 103645. <https://doi.org/10.1016/j.fm.2020.103645>.
- [28] Zhang R.; Gu J.; Wang X.; Li Y.; Liu J.; Lu C.; Qiu L. (2019). Response of antibiotic resistance genes abundance by graphene oxide during the anaerobic digestion of swine manure with copper pollution. *Science and Total Environment*, 654, 292–299. <https://doi.org/10.1016/j.scitotenv.2018.11.094>.
- [29] Zhang Y.; Zhou J.; Dong Z.; Li G.; Wang J.; Li Y.; Wan D.; Yang H.; Yin Y. (2019). Effect of dietary copper on intestinal microbiota and antimicrobial resistance profiles of *Escherichia coli* in weaned piglets. *Frontier in Microbiology*, 10, 1–11. <https://doi.org/10.3389/fmicb.2019.02808>.
- [30] USEPA (2007). Aquatic life ambient freshwater quality criteria - copper. R <https://www.epa.gov/wqc/aquatic-life-criteria-copper>.
- [31] Ong C.; Ibrahim S.; Sen Gupta B. (2007). A survey of tap water quality in Kuala Lumpur. *Urban Water Journal*, 4, 29–41. <https://doi.org/10.1080/15730620601145923>.
- [32] Del Olmo G.; Ahmad A.; Jensen H.; Karunakaran E.; Rosales E.; Calero Preciado C.; Gaskin P.; Douterelo I. (2020). Influence of phosphate dosing on biofilms development on lead in chlorinated drinking water bioreactors. *Biofilms and Microbiomes*, 6, 1–14. <https://doi.org/10.1038/s41522-020-00152-w>.
- [33] Wogu M.D.; Okaka C.E. (2011). Pollution studies on Nigerian rivers: Heavy metals in surface water of Warri river, Delta State. *Journal of Biodiversity and Environmental Science*, 1, 7–12.

- [34] Shanbehzadeh S.; Vahid Dastjerdi M.; Hassanzadeh A.; Kiyanzadeh T. (2014). Heavy metals in water and sediment: A case study of Tembi River. *Journal of Environmental and Public Health*, 2014, 858720. <https://doi.org/10.1155/2014/858720>.
- [35] Hadibarata, T.; Kristanti, R.A.; Mahmoud, A.H. (2020). Occurrence of endocrine-disrupting chemicals (EDCs) in river water and sediment of the Mahakam River. *Journal of Water and Health*, 18, 38-47, <https://doi.org/10.2166/wh.2019.100>.
- [36] Kaushik A.; Kansal A.; Santosh; Meena; Kumari S.; Kaushik C.P. (2009). Heavy metal contamination of river Yamuna, Haryana, India: Assessment by metal enrichment factor of the sediments. *Journal of Hazardous Materials*, 164, 265–270. <https://doi.org/10.1016/j.jhazmat.2008.08.031>.
- [37] Dippong T.; Mihali C.; Goga F.; Cical E. (2017). Seasonal evolution and depth variability of heavy metal concentrations in the water of Firiza-Strimtori Lake, NW of Romania. *Studia Universitatis Babeş-Bolyai Chemia*, 62, 213–228. <https://doi.org/10.24193/subbchem.2017.1.19>.
- [38] Purnaini R.; Sudarmadji; Purwono S. (2018). Tidal influence on water quality of Kapuas Kecil River downstream. E3S Web Conference, 31, 04006. <https://doi.org/10.1051/e3sconf/20183104006>.
- [39] Fu J.; Zhao C.; Luo Y.; Liu C.; Kyzas G.Z.; Luo Y.; Zhao D.; An S.; Zhu H. (2014). Heavy metals in surface sediments of the Jialu River, China: Their relations to environmental factors. *Journal of Hazardous Materials*, 270, 102–109. <https://doi.org/10.1016/j.jhazmat.2014.01.044>.
- [40] Bartoli G.; Papa S.; Sagnella E.; Fioretto A. (2012). Heavy metal content in sediments along the Calore river: Relationships with physical-chemical characteristics. *Journal of Environmental Management*, 95, S9–S14. <https://doi.org/10.1016/j.jenvman.2011.02.013>.
- [41] Kadhum S.A.; Ishak M.Y.; Zulkifli S.Z.; Hashim R.B. (2017). Investigating geochemical factors affecting heavy metal bioaccessibility in surface sediment from Bernam River, Malaysia. *Environmental Science and Pollution Research*, 24, 12991–13003. <https://doi.org/10.1007/s11356-017-8833-8>.
- [42] Zhang W.; Feng H.; Chang J.; Qu J.; Xie H.; Yu L. (2009). Heavy metal contamination in surface sediments of Yangtze River intertidal zone: An assessment from different indexes. *Environmental Pollution*, 157, 1533–1543. <https://doi.org/10.1016/j.envpol.2009.01.007>.
- [43] Li H.; Shi A.; Li M.; Zhang X. (2013). Effect of pH, temperature, dissolved oxygen, and flow rate of overlying water on heavy metals release from storm sewer sediments. *Journal of Chemistry*, 1–11. <https://doi.org/10.1155/2013/434012>.